

www.rockland.com tech@rockland.com +1 484.791.3823

#### Datasheet for 002-0303

# **Cat IgG Fc Peroxidase**

### **Overview**

Description:	Cat IgG Fc Fragment Peroxidase Conjugated - 002-0303
Item No.:	002-0303
Size:	1 mg
Origin:	Feline

### **Product Details**

Background:	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. This antibody possesses the F(c) region, recognized by high-affinity Fc receptor proteins.
Synonyms:	Cat IgG F(c) fragment HRP conjugated, Cat IgG F(c) fragment peroxidase conjugated
Species of Origin:	Feline
Conjugate:	Peroxidase (HRP)
Format:	IgG Fc
Туре:	Native Protein

### **Target Details**

**Purity/Specificity:** This product was prepared from normal serum by delipidation, salt fractionation, ion exchange

chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-

Peroxidase, anti-Cat IgG, anti-Cat IgG F(c) and anti-Cat Serum. No reaction was observed against

anti-Cat IgG F(ab')2 or anti-Papain.

## **Application Details**

**Application Note:** Cat IgG F(c) fragment Peroxidase conjugated can be utilized as a control or standard reagent in

Western Blotting and ELISA experiments.

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

Assay Dilutions:

All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

#### **Formulation**

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

### **Shipping & Handling**

<b>Shipping Condition:</b>	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

#### Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

www.rockland.com Page 3 of 3