

Datasheet for 200-401-E87**Amyloid Fibrils (OC) Antibody****Overview**

Description:	Anti-Amyloid Fibrils (OC) (RABBIT) Antibody - 200-401-E87
Item No.:	200-401-E87
Size:	100 µL
Applications:	IF, IHC, IP, WB, Other
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Amyloid monomeric proteins can sometimes oligomerize into destructive amyloid fibrils. Amyloidogenic conformations of non-disease related proteins can be created by partial protein misfolding or denaturation. Many degenerative diseases are known to be related to the accumulation of misfolded proteins as amyloid fibers. These include the amyloid- β peptide plaques and tau neurofibrillary tangles in senile plaques of Alzheimer's symptomology, the deposition of α -synuclein in the Lewy bodies of Parkinson's disease, and accumulation of polyglutamine-containing aggregates in Huntington's disease.
Synonyms:	Amyloid OC, Fibrils, Amyloid Oligomer $\alpha\beta$, A11, Amyloid beta A4 protein, ABPP, APPI, Alzheimer disease amyloid protein, Cerebral vascular amyloid peptide, PreA4, Protease nexin-II, APP, A4, AD1
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	APP
Reactivity:	Human
Immunogen Type:	Conjugated Peptide
Immunogen:	Amyloid Fibrils (OC) Antibody was produced from whole rabbit serum prepared by repeated immunizations with fibrils prepared from human A β 42 synthetic peptide.

Purity/Specificity:	Anti-Amyloid Fibrils (OC) Antibody was purified by Protein A chromatography. A BLAST analysis was used to suggest cross-reactivity with Amyloid Fibrils (OC) from Human based on 100% homology with the immunizing sequence. Expected to detect in mouse and rat based on species homology. Recognizes generic epitopes common to many amyloid fibrils and fibrillar oligomers, but not prefibrillar oligomers or natively folded proteins. Cross-reactivity with Amyloid Fibrils (OC) from other sources has not been determined. Neuroscience research.
----------------------------	---

Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P05067
------------------------	--

Application Details

Tested Applications:	IF, IHC, IP, WB
Suggested Applications:	Other (Based on references)
Application Note:	Anti-Amyloid Fibrils (OC) Antibody is tested for use in IP, IF microscopy, IHC, and WB. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:200
IHC:	User Optimized
IP:	User Optimized
WB:	1:1000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	N/A by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.09% (w/v) Sodium Azide
Stabilizer:	50% (v/v) Glycerol


Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images




Immunohistochemistry

Immunohistochemistry of Rabbit anti-Amyloid Fibrils antibody. Tissue: Human AD Brain. Fixation: N/A. Primary Antibody: (Left) Amyloid Fibril antibody, (Right) Monoclonal 6E10 at 1ug/ml for 1h at RT. Secondary antibody: Peroxidase rabbit secondary at 1:10,000 for 45 min at RT. Localization: Membrane. Staining: (Left) Amyloid Fibrils as precipitated brown signal with no cross reactivity with Amyloid Precursor Protein (APP), (right) considerable cross reactivity.




Immunohistochemistry

Immunohistochemistry of Rabbit anti-Amyloid Fibrils antibody. Tissue: (A) hippocampus, (B) subiculum, (C) temporal cortex, and (D) dense and fine fibrillar material. Fixation: N/A. Primary Antibody: Amyloid Fibrils antibody at 1ug/ml for 1h at RT. Secondary antibody: Peroxidase rabbit secondary at 1:10,000 for 45 min at RT. Localization: Membrane. Staining: Amyloid Fibrils as precipitated brown signal with hematoxylin purple nuclear counterstain.




Dot Blot

Dot Blot of Rabbit Amyloid Fibrils (OC) antibody. Antigen: Aβ42 and polyQ36 prefibrillar oligomers and fibrils. Load: 2ug per dot. Primary antibody: Top row: Amyloid Oligomers (A11) or bottom row: Amyloid Fibrils (OC) at 1:400 for 45 min at 4°C. Secondary Antibody: Goat anti-rabbit IgG HRP at 1:10,000 for 45 min at RT. Block: 5% Blotto overnight at 4°C. Amyloid Fibrils (OC) reacts to Aβ42 fibrils and polyQ36 fibrils only.



Dot Blot

Dot Blot of Rabbit anti-Amyloid Fibrils antibody. Antigen: Beta Amyloid HEPES-NaCl aggregation. Primary antibody: Amyloid Fibrils antibody at 1:500 (lane 1) and 1:5000 (lane 2) for 45 min at 4°C. Secondary antibody: Peroxidase rabbit Secondary antibody at 1:10,000 time lapse. Block: 5% BLOTTO overnight at 4°C.



Western Blot

Western Blot of rabbit Anti-Amyloid Fibrils Antibody. Lane 1 and 3: (F) Fibrils. Lane 2 and 4: (O) prefibrillar oligomers. Load: 10 ug per lane. Primary antibody: Anti-Amyloid Fibrils or Anti-Oligomers at 1:1000 for overnight at 4°C. Secondary antibody: Goat anti-rabbit IgG HRP antibody at 1:40,000 for 45 min at RT. Block: 5% Blotto overnight at 4°C. Predicted/Observed size: 18kDa on left blot (OC) in lane one.

References

- Lee D. et al. Plasmonic nanoparticle amyloid corona for screening A β oligomeric aggregate-degrading drugs. *Nat Commun.* (2021)
- Nnah IC et al. Iron potentiates microglial interleukin-1 β secretion induced by amyloid- β . *J Neurochem.* (2020)
- Jeong D et al. Multifunctionalized Reduced Graphene Oxide Biosensors for Simultaneous Monitoring of Structural Changes in Amyloid- β 40. *Sensors (Basel).* (2018)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.