

www.rockland.com tech@rockland.com +1 484.791.3823

#### Datasheet for 600-401-J72

# **HDAC8** (near N-terminus) Antibody

### **Overview**

Description:	Anti-HDAC8 (RABBIT) Antibody - 600-401-J72
Item No.:	600-401-J72
Size:	100 μg
Applications:	ELISA, WB
Reactivity:	Human
Host Species:	Rabbit

### **Product Details**

Background:	HDAC8 is located in the nucleus and cytoplasm, expressed weakly in most tissues with higher expression in the heart, brain, kidney, pancreas, liver, lunch, placenta, prostate, and kidney. It is responsible for the deacetylation of lysine residues on the N-terminal region of the core histones (H2A, H2B, H3 and H4). The result of deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Anti-HDAC8 therefore is ideal for investigators interested in Stem Cells , Cardiovascular , or Epigenetics and Nuclear Signaling research.
Synonyms:	rabbit anti-HDAC-8 antibody, HDAC8, HDAC 8, HD-8, HD 8, Histone deacetylase 8, Histone deacetylase-8, HD8, HDACL1, CDA07
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

## **Target Details**

Gene Name:	HDAC8
Reactivity:	Human
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-HDAC8 affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide near the N-terminus of human HDAC8 protein.

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

Purity/Specificity: Anti-HDAC8 was affinity purified from monospecific antiserum by immunoaffinity

chromatography. A BLAST analysis was used to suggest cross-reactivity with rat, mouse, human, and bovine based on 100% sequence homology. Cross-reactivity with HDAC8 from

other sources has not been determined.

Relevant Links: • UniProtKB - Q9BY41

### **Application Details**

<b>Tested Applications:</b>	ELISA, WB
Application Note:	Anti-HDAC8 antibody has been tested by ELISA, and western blot and is useful for Immunohistochemistry. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~41kDa corresponding to the appropriate cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1: 10,000
WB:	1ug/ml

### **Formulation**

Physical State:	Liquid (sterile filtered)
Concentration:	1.0mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer:	30% Glycerol

### **Shipping & Handling**

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

#### Disclaimer

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3