

#### Datasheet for 605-744-125

# Goat IgG (H&L) Antibody DyLight™ 680 Conjugated Pre-Adsorbed

## **Overview**

Description:	Donkey Anti-Goat IgG (H&L) Antibody DyLight™ 680 Conjugated (Min X Ch GP Ham Hs Ms Rb & Rt Serum Proteins) - 605-744-125
Item No.:	605-744-125
Size:	100 μg
Applications:	WB
Reactivity:	Goat
<b>Host Species:</b>	Donkey

## **Product Details**

Background:	Anti-Goat IgG DyLight Anti	body genera	ated in donkey	detects goat IgG.	Secreted as part of the
	the state of the s				

adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F (ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody

molecule are present.

Synonyms: Donkey anti-Goat IgG Antibody DyLight™ 680 Conjugated Pre-Adsorbed, Donkey anti-Goat IgG

DyLight™ 680 Conjugated Antibody

Host Species: Donkey

**Specificity:** IgG (H&L)

Conjugate: DyLight™ 680

**Clonality:** Polyclonal

Format: IgG

**F/P Ratio:** 3.4

# **Target Details**

Reactivity: Goat

www.rockland.com Page 1 of 4





Immunogen:	Goat IgG whole molecule
Purity/Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Goat IgG and Goat Serum. No reaction was observed against Chicken, Guinea Pig, Hamster, Horse, Mouse, Rabbit and Rat Serum Proteins. This antibody will react with heavy chains of Goat IgG and with light chains of most Goat immunoglobulins.

# **Application Details**

<b>Tested Applications:</b>	WB
Application Note:	Anti-Goat IgG DyLight 680 Antibody has been tested by western blot. This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. The emission spectra for this DyLight™ conjugate match the principle output wavelengths of most common fluorescence instrumentation.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
FLISA:	>1:20,000
IF:	>1:5,000
WB:	>1:10,000

# **Formulation**

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	100 μL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

# **Shipping & Handling**

Shipping Condition: Ambient

www.rockland.com Page 2 of 4






**Storage Condition:** Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20°

C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an

undiluted liquid. Dilute only prior to immediate use.

**Expiration:** Expiration date is one (1) year from date of receipt.

# **Images**



#### **Western Blot**

DyLight™ dyes can be used for two-color Western Blot detection with low background and high signal. Anti-tubulin was detected using a DyLight™ 680 conjugate. Anti-TNFa was detected using a DyLight™ 800 conjugate. The image was captured using the Odyssey® Infrared Imaging System developed by LI-COR.

## Diagram


Properties of DyLight™ Fluorescent Dyes

Emission	Color	DyLight™ Dye	Ex/Em (nm)	e (M⁻¹ cm⁻¹)	Similar Dyes
Blue		405	400/420	30,000	Alexa™ 405, Cascade Blue
Green		488	493/518	70,000	Alexa <sup>™</sup> 488, Cy2 <sup>®</sup> , FITC
Yellow		549	550/568	150,000	Alexa™ 546, Alexa 555, Cy3®,TRITC
Red		649	646/674	250,000	Alexa™ 647, Cy5®
Near Infrared	7	680	682/715	140,000	Alexa™ 680, Cy5.5®, IRDye™ 700
Infrared		800	770/794	270,000	IRDye™ 800

www.rockland.com Page 3 of 4








#### **Western Blot**

Western Blot of Donkey anti-Goat IgG Pre-Adsorbed DyLight™ 680 Conjugated Secondary Antibody. Lane 1: Goat IgG. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: DyLight™ 680 goat secondary antibody at 1:5,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Goat IgG. Other band(s): None.



# Diagram

# References

• Van den Hoecke S et al. In Vivo Therapy with M2e-Specific IgG Selects for an Influenza A Virus Mutant with Delayed Matrix Protein 2 Expression. *mBio*. (2021)

## **Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 4 of 4