

www.rockland.com tech@rockland.com +1 484.791.3823

#### **Datasheet for KOA0214**

### **Human IL-2 ELISA Kit**

#### **Overview**

Description:	Human IL-2 AccuSignal ELISA Kit - KOA0214
Item No.:	KOA0214
Size:	1 Kit
Applications:	ELISA
Reactivity:	Human
Range:	15.6 pg/ml - 1000 pg/ml

#### **Product Details**

**Background:** Interleukin-2(IL2), formerly referred to as T-cell growth factor, is a powerfully immunoregulatory

lymphokine that is produced by lectin- or antigen-activated T cells. It is produced not only by mature T lymphocytes on stimulation but also constitutively by certain T-cell lymphoma cell lines. The lymphokine interleukin-2(IL-2) is responsible for autocrine cell cycle progression and regulation of immune responses. IL-2 expression in mature thymocytes and T cells has been found to be tightly controlled by monoallelic expression. IL-2 can act as a growth hormone for both B and T lymphocytes. The human gene for interleukin 2(IL2) is assigned to chromosome 4. Human IL-2 is a 15.5KDa glycoprotein, consisting of 153 amino acids in precursor form and 133

amino acids in mature form.

Synonyms: Aldesleukin, IL 2, IL-2, IL2\_HUMAN, Interleukin 2, Interleukin-2, Interleukin-2, Involved in

regulation of T cell clonal expansion, Lymphokine, T cell growth factor, T-cell growth factor,

**TCGF** 

**Detection Kit Type:** ELISA Kit

**Detection Range:** 15.6 pg/ml - 1000 pg/ml

**Sensitivity:** <1 pg/ml

## **Target Details**

Gene Name:	IL2
Reactivity:	Human
Immunogen:	Expression system for standard: E.coli; Immunogen sequence: A21-T153

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

Purity/Specificity:	Natural and recombinant human IL-2. There is no detectable cross-reactivity with other relevant proteins.
Relevant Links:	• UniProtKB - P60568
	• NCBI - NP_000577.2
	• GeneID - 3558
	KOA0214 Protocol

# **Application Details**

Tested Applications:	ELISA
Application Note:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 1000pg/ml, 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml, 31.2pg/ml, 15.6pg/ml human IL-2 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of human cell culture supernates, serum or plasma( heparin, EDTA, citrate) to each empty well. It is recommended that each human IL-2 standard solution and each sample be measured in duplicate.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	DOE: 2-8° C - 02/26/2020 -20° C - 08/26/2020

### **Formulation**

Concentration:	1 kit
Anticoagulant:	Heparin Sodium

# **Shipping & Handling**


Shipping Condition:	Wet Ice
Storage Condition:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
Expiration:	See kit insert for complete instructions.

www.rockland.com Page 2 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

### **Images**



#### **ELISA**

Human IL-2 AccuSignal ELISA Kit standard curve. Assay Range: 15.6pg/ml-1000pg/ml. Sensitivity: <1pg/ml.

#### **Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3