

www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for KOA0770

Mouse Angiopoietin-1 ELISA Kit

Overview

Description:	Mouse Angiopoietin-1 AccuSignal ELISA Kit - KOA0770
Item No.:	KOA0770
Size:	1 Kit
Applications:	ELISA
Reactivity:	Mouse
Range:	156 pg/ml - 10,000 pg/ml

Product Details

Background: Angiopoietin 1, also called ANG1 is a type of angiopoietin and is encoded by the gene ANGPT1.

Angiopoietins are proteins with important roles in vascular development and angiogenesis. All angiopoietins bind with similar affinity to an endothelial cell-specific tyrosine-protein kinase receptor. This gene was mapped to 8q23.1. The protein encoded by this gene is a secreted glycoprotein that activates the receptor by inducing its tyrosine phosphorylation. It plays a critical role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme and inhibits endothelial permeability. The protein also contributes to blood vessel maturation and stability, and may be involved in early development of the heart.

Synonyms: 1110046O21Rik, AGP 1, AGP1, AGP1, ANG 1, ANG1, Angiopoietin1, ANGPT 1, ANGPT1, KIAA0004

Detection Kit Type: ELISA Kit

Detection Range: 156 pg/ml - 10,000 pg/ml

Sensitivity: <10 pg/ml

Target Details

Gene Name:	ANGPT1
Reactivity:	Mouse
Immunogen:	Expression system for standard: NSO; Immunogen sequence: H16-F497
Purity/Specificity:	Natural and recombinant mouse Angiopoietin-1. There is no detectable cross-reactivity with other relevant proteins.

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

Relevant Links: • UniProtKB - 008538

• NCBI - NP_033770.2

• GeneID - 11600

KOA0770 Protocol

Application Details

Tested Applications:	ELISA
Application Note:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 10,000pg/ml, 5000pg/ml, 2500pg/ml, 1250pg/ml, 625pg/ml, 312pg/ml, 156pg/ml mouse Angiopoietin-1 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended that each mouse Angiopoietin-1 standard solution and each sample be measured in duplicate.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

Formulation

Anticoagulant: Heparin Sodium

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
Expiration:	See kit insert for complete instructions.


Images

www.rockland.com Page 2 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

Mouse Angiopoietin-1 ELISA Kit



ELISA

Mouse Angiopoietin-1 ELISA Kit standard curve.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3